

MA2219

THE UNIVERSITY OF WARWICK

SECOND YEAR RESIT EXAMINATION: JUNE 1996

GEOMETRY AND TOPOLOGY

Time allowed: 2 hours

Read carefully the instructions on the answer book and make sure that the particulars required are entered on it.

ANSWER 4 QUESTIONS

If you have answered more than the required 4 questions, you will only be given credit for your 4 best answers.

1. (i) What is meant by a motion of Euclidean n -space $\mathbf{E}^n$? State and prove a necessary and sufficient condition on an $n \times n$ matrix A in order that the map defined by

$$\mathbf{x} \mapsto A\mathbf{x} + \mathbf{b} \quad \text{for } \mathbf{x} \in \mathbf{E}^n \quad (*)$$

should be a motion of $\mathbf{E}^n$.

- (ii) Let L_1 and L_2 be the lines of $\mathbf{E}^2$ defined by the equations

$$L_1 : (y = b) \quad \text{and} \quad L_2 : (\cos \theta)y = (\sin \theta)x,$$

where θ is a fixed angle between 0 and π , and $\beta \in \mathbf{R}$. Write down the reflections R_1, R_2 in L_1 and L_2 in the form $(*)$; proofs are not required.

- (iii) Calculate the composite motion $R_2 \circ R_1$ in this form.
(iv) Using the fact that both R_1 and R_2 fix the point

$$P = L_1 \cap L_2 = ((\cot \theta)\beta, \beta),$$

describe the effect of $R_2 \circ R_1$ on $\mathbf{E}^2$ in geometric terms.

CONTINUED ...

2. (a) Define the notions of affine linear subspace and linear span in affine space $\mathbf{A}_{\mathbf{R}}^n$. State and prove the theorem on dimension of intersection of affine linear subspaces $E, F \subset \mathbf{A}_{\mathbf{R}}^n$ (results on vector spaces may be used freely). Explain briefly why this result is more convenient to state in projective space $\mathbf{P}_{\mathbf{R}}^n$.
- (b) Consider $\triangle PQR$ and $\triangle P'Q'R'$ in projective 3-space $\mathbf{P}^3$, where

$$P = (0, 0, 0, 1), \quad Q = (1, 0, 0, 1), \quad R = (0, 1, 0, 1),$$

$$P' = (0, 0, 1, 1), \quad Q' = (1, 0, 1, 1), \quad R' = (0, 1, 1, 1).$$

By writing out the lines PP' , QQ' , RR' , show that $\triangle PQR$ and $\triangle P'Q'R'$ are in perspectivity. [Hint: you may want to take coordinates x, y, z, t and to visualise $PQRP'Q'R'$ as a triangular prism in the affine piece $t = 1$.] Now determine the points of intersection of the corresponding sides of the triangles, for example:

$$QR : (x + y = t, z = 0), \quad Q'R' : (x + y = t, z = t),$$

$$\implies QR \cap Q'R' = (1 : -1 : 0 : 0).$$

- (c) State and prove Desargues' theorem for two triangles $\triangle PQR$ and $\triangle P'Q'R'$ spanning distinct planes in $\mathbf{P}_{\mathbf{R}}^3$ and in perspectivity from a point O . Verify it in the example of (b).

CONTINUED ...

3. (i) Define a reflection of Euclidean n -space $\mathbf{E}^n$, and state without proof the theorem of the course expressing any motion of $\mathbf{E}^n$ as a composite of a number of reflections. Explain what it means in the particular case of a direct motion of $\mathbf{R}^3$ fixing the origin. Deduce that a 3×3 orthogonal matrix A with $\det A = 1$ has $+1$ as eigenvalue. (That is, any element $A \in \text{SO}(3)$ has a fixed axis.)
- (ii) Define the quaternion conjugate q^* of a quaternion q , and explain how to find the inverse q^{-1} if $q \neq 0$. Let p, q be quaternions. Prove that

$$p \text{ is pure imaginary} \implies qpq^{-1} \text{ is pure imaginary.}$$

Identify the space of pure imaginary quaternions $p = xi + yj + zk$ with Euclidean space $\mathbf{E}^3$ (with coordinates x, y, z). Assuming that $|q|^2 = 1$ for simplicity, prove that the map $r_q: \mathbf{E}^3 \rightarrow \mathbf{E}^3$ given by $r_q(p) = qpq^{-1}$ is an element of $\text{SO}(3)$.

- (iii) Determine the rotation r_q for the unit quaternion

$$q = c + si \quad \text{with} \quad c^2 + s^2 = 1,$$

and express the result geometrically.

4. (a) Let X be a topological space and $Y \subset X$ a subset with the subspace topology. Under the assumption that X is compact, prove that Y closed implies Y compact. Under what condition on X is the converse true? (The proof is not required).
- (b) Let X and Y be compact metric spaces and $\varphi: X \rightarrow Y$ a continuous bijective map. Sketch the proof that φ is a homeomorphism.
- (c) Let $S^1 = \{(x, y) \in \mathbf{R}^2 \mid x^2 + y^2 = 1\}$ be the unit circle, and $\varphi: S^1 \rightarrow \mathbf{R}^n$ a continuous map with the property that for all $(x, y) \in S^1$

$$\varphi(x, y) = \varphi(x', y') \iff (x, y) = \pm(x', y').$$

Prove that the image $\varphi(S^1)$ is homeomorphic to S^1 .

CONTINUED ...

5. Use your mathematical judgement to argue for the truth or otherwise of the following statements.
- (a) Let S_n be the symmetric group on $\{1, 2, \dots, n\}$ and $G \subset S_n$ a subgroup; if G contains the transposition (12) and an element taking $i \mapsto 1, j \mapsto 2$, for any pair i, j of distinct elements of $\{1, 2, \dots, n\}$, then $G = S_n$.
 - (b) There exists a finite non-Abelian group of direct motions of the Euclidean plane $\mathbf{E}^2$.
 - (c) There exists an affine linear map $\mathbf{A}^1 \rightarrow \mathbf{A}^1$ of the affine line taking any set of 3 distinct points into any other.
 - (d) Given 3 lines $L_1, L_2, L_3 \subset \mathbf{P}^3$, there is a unique line M meeting all 3.
 - (e) Let (X, d) be a metric space and $\sim$ an equivalence relation on X ; then setting

$$d([x], [y]) = \inf \{d(x', y')\} \quad \text{taken over all } x' \in [x], y' \in [y].$$

defines a metric on the quotient space $Y = X / \sim$.

END

Answer to Q1.

(i) A motion of $\mathbf{E}^n$ is a map $t: \mathbf{E}^n \rightarrow \mathbf{E}^n$ that preserves Euclidean distances, i.e., $d(t(P), t(Q)) = d(P, Q)$ for all P, Q .

[Definition, 3 marks]

If t is given by $x \mapsto Ax + b$ with A and $n \times n$ matrix and b a vector, t is a motion if and only if A is orthogonal. Indeed, the translation by b does not affect distances, so that I need to prove that $x \mapsto Ax$ preserves distance from the origin, that is, the quadratic form $|x|^2 = {}^t x \cdot x$. But this holds if and only if it preserves the bilinear form ${}^t x x$ on $\mathbf{R}^n$, that is,

$${}^t(Ax) \cdot Ay = {}^t x ({}^t A \cdot A)y = {}^t x y \quad \text{for all } x, y \in \mathbf{R}^n,$$

that is, ${}^t A A = 1$.

[Bookwork, 6 marks]

(ii) The reflection in $L_1 : (y = \beta)$ is given by $x \mapsto x, y \mapsto 2\beta - y$, that is,

$$R_1: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ 2\beta \end{pmatrix}.$$

The reflection in $L_2 : (\cos \theta)y = (\sin \theta)x$ is given by

$$R_2: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

(Several derivations are possible, for example, by computing the conjugate $\text{Rotation}(\theta) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \text{Rotation}(-\theta)$, or by finding the fixed axis.)

[4+4 marks]

(iii) So

$$\begin{aligned} R_2 \circ R_1: \begin{pmatrix} x \\ y \end{pmatrix} &\mapsto \begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix} \left[\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ 2\beta \end{pmatrix} \right] \\ &= \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + 2\beta \begin{pmatrix} \sin 2\theta \\ -\cos 2\theta \end{pmatrix} \end{aligned}$$

[4 marks]

(iv) $R_2 \circ R_1 = \text{Rotation}(P, 2\theta)$, where as given, $P = L_1 \cap L_2 = ((\cot \theta)\beta, \beta)$ (see Figure).

[4 marks]

[Standard type of question. $3 + 6 + 4 + 4 + 4 + 4 = 25$]

Answer to Q2.

(a) $\mathbf{A}^n = \mathbf{R}^n$. An affine linear subspace is of the form $u_0 + V$ where $u_0 \in \mathbf{R}^n$ and $V \subset \mathbf{R}^n$ is a vector subspace. Given a nonempty set $\Sigma \subset \mathbf{A}^n$, its span $\langle \Sigma \rangle$ is the smallest affine linear subspace containing Σ . In other words, if $u_0 \in \Sigma$ is any point,

$$\langle \Sigma \rangle = u_0 + V, \quad \text{where} \quad V = \langle \Sigma - u_0 \rangle.$$

The theorem says that for E, F affine linear subspaces,

$$\dim E + \dim F = \dim(E \cap F) + \dim \langle E, F \rangle,$$

provided $E \cap F \neq \emptyset$. For the proof, just pick $u_0 \in E \cap F$, so that $E = u_0 + U$, $F = u_0 + V$ for some vector subspaces $U, V \subset \mathbf{R}^n$. Then

$$E \cap F = u_0 + (U \cap V), \quad \langle E, F \rangle = u_0 + (U + V),$$

and the formula therefore follows from that for vector spaces.

The same formula holds for all E, F without exception in $\mathbf{P}^n$, provided we write $\dim \emptyset = -1$. The lecturer seemed to think this was really good.

[Bookwork. 11 marks]

(b) $PP' : (x = y = 0)$, $QQ' : (x = t, y = 0)$ and $RR' : (x = 0, y = t)$ intersect in $O = (0010)$. (By the hint, the sides of the prism are vertical and parallel so meet at (0010) .)

Now $PQ : (y = z = 0)$, $P'Q' : (y = 0, z = t)$ intersect in $C = (1000)$,

$PR : (x = z = 0)$, $P'R' : (x = 0, z = t)$ intersect in $B = (0100)$,

and $QR \cap Q'R'$ intersect in $A = (1 : -1 : 0 : 0)$ is given.

[Easier than lectured material. 7 marks]

(c)

Desargues' theorem. *If $\triangle PQR$ and $\triangle P'Q'R'$ are in perspectivity from a point O then the 3 points of intersection*

$$A = QR \cap Q'R', \quad B = PR \cap P'R', \quad C = PQ \cap P'Q'$$

of the corresponding sides are collinear.

Proof. If the two triangles span distinct planes $\Pi = \langle \triangle PQR \rangle \neq \Pi' = \langle \triangle P'Q'R' \rangle$ then this is obvious: $OPQP'Q'$ is a plane so PQ and $P'Q'$ intersect in a point C , etc., and so C is contained in lines in the planes Π , Π' , therefore is contained in the line $L = \Pi \cap \Pi'$. Similarly for B, C . Q.E.D.

The above 3 points $(1 : -1 : 0 : 0), (0100), (1000)$ are obviously on $z = t = 0$.

[The easy case of Desargues theorem. 3 marks. $11 + 7 + 7 = 25$]

Answer to Q3.

(i) The reflection of $\mathbf{E}^n$ in a hyperplane Π is the motion that fixes every $P \in \Pi$, and that takes a point Q into $t(Q)$ constructed as the mirror image, that is, drop a perpendicular from Q to Π and set $t(Q)$ to be the point the same distance the other side. (In other words, Π is the perpendicular bisector of PQ .)

Theorem. *Any motion of $\mathbf{E}^n$ is a composite of at most $n + 1$ reflections. Any motion fixing the first k elements of a Euclidean frame of reference is a composite of at most $n + 1 - k$ reflections.*

In particular, if $n = 3$ and $k = 1$, any motion $t: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ is a composite of ≤ 3 reflections. If t is direct then the number of reflections is even, that is, 0 (the identity) or 2. But the composite of 2 distinct reflections $\text{Reflection}(\Pi_1), \text{Reflection}(\Pi_2)$ fixes the line $L = \Pi_1 \cap \Pi_2$ (vector subspaces), as required.

[Some students will probably do the last part “Otherwise” for fewer marks: There exists an orthonormal basis w.r.t. which A is the block diagonal matrix, with each block either 1×1 block with entry ± 1 , or the 2×2 rotation matrix

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

In the 3×3 case there is at most one rotation block. If $\det A = 1$ (that is, A is direct) then either $A = 1_3$ or A has a block $+1$ and a rotation block.]

[Statement of theorem. Deconstructing to $\mathbf{R}^3$ is unseen. 6+4 marks]

(ii) If $q = a + bi + cj + dk$ then $q^* = a - bi - cj - dk$. Thus q is real resp. pure imaginary if $q^* = q$ or $q^* = -q$. If $q \neq 0$ then

$$qq^* = |q|^2 = a^2 + b^2 + c^2 + d^2$$

is a positive real, so that $q^{-1} = q^*/|q|^2$ is the inverse.

Now for a real, $qaq^* = aqq^* = a|q|^2$ is also real. If p is pure imaginary then $p^* = -p$, so that $(qpq^*)^* = qp^*q^* = -qpq^*$, and p is imaginary.

Setting $r_q(p) = qpq^{-1}$ then r_q is a linear map $\mathbf{R}^3 \rightarrow \mathbf{R}^3$, and $|r_q(p)|^2 = |qpq^{-1}|^2 = (qpq^*)(qp^*q)/|q|^4 = pp^* = |p|^2$, so that r_q is a motion or orthogonal map. It is direct because $\det r_q$ is a continuous function from the nonzero quaternions to ± 1 .

[Bookwork. 7 marks]

(d) Write $c = \cos \psi$, $s = \sin \psi$ and $q = c + si$. Then q is a unit quaternion; $r_q(i) = qiq^{-1} = i$ because $\mathbf{R}[i] = \mathbf{C}$ is commutative. And the action on j, k is given by

$$\begin{aligned} r_q(j) &= (c + si)j(c - si) = (c^2 - s^2)j + 2csk \\ \text{and} \quad r_q(k) &= (c + si)k(c - si) = -2csj + (c^2 - s^2)k. \end{aligned}$$

This is a rotation of the y, z plane by an angle 2ψ .

[Calculation similar to proof in course. 8 marks. $10 + 7 + 8 = 25$]

Answer to Q4.

(a) X is compact and $Y \subset X$ closed. Let U_λ be an open cover of Y indexed by $\lambda \in \Lambda$. By definition of the subspace topology, there exist open sets $V_\lambda \subset X$ with $U_\lambda = V_\lambda \cap Y$. Then $Y \subset \bigcup V_\lambda$, so that $X = CY \cup \bigcup V_\lambda$ is an open cover of X (where CY is the complement of Y). Since X is compact, a finite number of these cover X , say $X = CY \cup V_i$, so $Y \subset \bigcup V_i$, that is, $Y = \bigcup (V_i \cap Y) = \bigcup U_i$. This proves Y compact.

[Bookwork. 7 marks]

The converse is true if X is Hausdorff: this means for all $x, y \in X$, there exist disjoint open sets U_x, U_y with $x \in U_x, y \in U_y$.

[Bookwork, I'll buy metric instead. 3 marks]

(b) φ is bijective, so that there is an inverse map φ^{-1} . If we prove that φ is a closed map (that is, takes closed sets of X to closed sets of Y) then φ^{-1} is continuous, so that φ is a homeomorphism.

Now X and Y are compact and Hausdorff by assumption, so that closed = compact by (a). Now a continuous map φ takes compact sets to compact sets, therefore it takes closed sets to closed sets.

[Bookwork. 6 marks]

(c) Let $\sim$ be the equivalence relation $(x, y) \sim \pm(x, y)$. Then by assumption, φ is constant on the equivalence classes, so defines a continuous injective map $\psi: S^1/(\pm 1) \rightarrow \mathbf{R}^n$. This map is a homeomorphism of $S^1/(\pm 1)$ to its image $\psi(S^1)$ by the theorem of (b).

The point is to see that $S^1/(\pm 1) \simeq S^1$, for example because it is the upper half-circle with its ends identified.

[Similar to $SO(3) = S^3/\pm 1$ in course. Bit long, but making it shorter would make it harder. (c) contains a trap for the unwary. 9 marks.]

[7 + 3 + 6 + 9 = 25]

Answer to Q5.

(a) True. By the given information, G contains (12) , and $g = g_{ij} : i \mapsto 1, j \mapsto 2$, so by the conjugacy principle, it contains $g(12)g^{-1} = (ij)$. Obviously S_n is generated by all the transpositions.

[Tests conjugacy, 4.2–3 of book. 5 Marks]

(b) False. A finite group $G \subset \text{Eucl}(2)$ fixes a point P_0 (the centroid of any orbit $G \cdot P$); therefore G has only reflections and rotations. If it's direct it only contains rotations around P_0 , therefore is commutative. Thus the statement is false.

[Tests various aspects of Euclidean groups. 5 Marks]

(c) False. The ratio of distances is defined. The affine linear map $\varphi: \mathbf{A}^1 \rightarrow \mathbf{A}^1$ given by $x \mapsto ax + b$ is determined by its effect on $P = 0$ and $Q = 1$: because $\varphi(0) = b$, and $\varphi(1) = a + b$. Therefore, given another point $R = c$, the image $\varphi(R) = ac + b$ is determined by $\varphi(P)$ and $\varphi(Q)$, so can't be changed.

[Easy but unseen; like cross-ratio. 5 Marks]

(d) False. There are infinitely many. For a counterexample, if L_1 and L_2 are disjoint, they span $\mathbf{P}^3$ (by dimension of intersection), so every point of L_3 is on a line joining L_1, L_2 , and this make infinitely many lines

[Simple trap. 5 Marks]

(e) As false as can be: the inf can be zero. For example, in $\mathbf{R}$, set $x \sim y$ iff either both nonzero or both zero. Then $x \not\sim y$ means $d(x, y) > 0$ but not bounded below.

Also, the triangle inequality does not hold: for example, if $\sim$ identifies the ends of $[0, 1]$ (to give the circle), if $x, z =$ points near the cuts and $y = 0 \sim 1$, then $d(x, [y]), d([y], z)$ are very small, but $d(x, z)$ is near to 1.

[Copied from past exam made available to students]

[By the warning, ≤ 1 mark for true/false. $5 \times 5 = 25$]